

4300430: Introdução à Cosmologia Física

Problem Set 9

(Due November 07, 2024)

1) Correlation Function in Redshift Space. In class, we saw that the correlation function in redshift space is anisotropic in space and given by

$$\xi_g^s(\mathbf{s}) = \xi_g^s(s_{\parallel}, s_{\perp}) = \xi_g^s(s, \mu_s) = b^2 \sum_{l=0,2,4} c_l(\beta) L_l(\mu_s) \xi_l^s(s) \quad (1)$$

where $L_l(\mu_s)$ is the Legendre Polynomial of order l , $\mu_s = \cos(\theta_s)$ is the cosine of the angle between the vector \mathbf{s} and the line-of-sight $\hat{\mathbf{z}}$, the coefficients

$$c_l(\beta) = \frac{2l+1}{2} \int_{-1}^1 (1+\beta x^2)^2 L_l(x) dx = \begin{cases} 1 + \frac{2}{3}\beta + \frac{1}{5}\beta^2, & l=0 \\ \frac{4}{3}\beta + \frac{4}{7}\beta^2, & l=2 \\ \frac{8}{35}\beta^2, & l=4 \end{cases} \quad (2)$$

where $\beta = f/b$, b is the galaxy bias and $f = \frac{d \ln D}{d \ln a}$, and the multipoles

$$\xi_l^s(s) = i^l \int \frac{k^2 dk}{2\pi^2} j_l(ks) P^r(k) \quad (3)$$

Assume the fiducial cosmology from previous problem sets in the calculations below.

a) From the real-space matter power spectrum $P^r(k)$ (e.g. from CAMB), use Eq.(3) to compute the multipoles $\xi_l^s(s)$ for $l = 0, 2, 4$. Plot each multipole as a function of separation s in log-scale and appropriate ranges. Notice that if the spectrum has k in units of h/Mpc , the separation s will naturally be in units of Mpc/h . Similarly $P^r(k)$ is in units of $[\text{Mpc}/h]^3$, and therefore the multipoles are unitless. Note that $j_0(x) = \sin(x)/x$ and other values of l can be obtained by recurrence relations.

b) Assuming $b = 1$, compute explicitly the analytical integral in Eq.(2) to derive the coefficients $c_l(\beta)$. If you have a numerical growth function $D(z)$ from the previous problem set 8, use it to compute f ; otherwise use a fitting function [e.g. $f = \Omega_m^\gamma(z)$ where $\Omega_m(z) = \Omega_m(1+z)^3/E^2(z)$]. Then plot $f(z)$ and $c_l[\beta(z)]$ (for $l = 0, 2, 4$) as a function of redshift z .

c) Finally, use Eq.(1) to obtain the redshift-space correlation function as a function of parallel and perpendicular directions at $z = 0$. Make a 2D plot of your results, with $s_\perp = s \sin \theta$ in the x-axis, $s_\parallel = s \cos \theta_s$ in the y-axis, and a color coding for the value of $\xi^s(s_\perp, s_\parallel)$. Compare these results to a similar 2D plot for the isotropic real-space correlation $\xi^r(s) = \xi_{l=0}^s(s)$. Repeat the same plots at $z = 1.0$.