

PGF5292: Physical Cosmology I

Problem Set 11

(Due June 03, 2015)

1) Halo Model Ingredients

In PS4, you computed the halo mass-function for a fit from Tinker et al. 2008

$$\frac{dn(z, M_{vir})}{d \ln M_{vir}} = f(\sigma) \frac{\bar{\rho}_{m0}}{M_{vir}} \frac{d \ln \sigma^{-1}}{d \ln M_{vir}} \quad (1)$$

$$f(\sigma) = A \left[\left(\frac{\sigma}{b} \right)^{-a} + 1 \right] \exp \left[-\frac{c}{\sigma^2} \right] \quad (2)$$

Now you must set values for A, a, b, c appropriate for $\Delta = \Delta_c$ (Eq. 11 below). Here $\sigma^2(M_{vir})$

$$\sigma^2(z, R) = D^2(z) \int \frac{k^2 dk}{2\pi^2} |W(kR)|^2 P_L(k) \quad (3)$$

where $M = \bar{\rho}_{m0} 4\pi R^3 / 3$, $P_L(k)$ is the linear power spectrum at redshift $z = 0$ and

$$W(kR) = \frac{3}{k^2 R^2} \left[\frac{\sin(kR)}{kR} - \cos(kR) \right] \quad (4)$$

is the Fourier Transform of a spherical top-hat window of radius R .

a) For a fiducial cosmology, compute the halo bias for the fit from Tinker et al. 2010:

$$b(z, M) = 1 - A \frac{\nu^a}{\nu^a + \delta_c^a} + B\nu^b + C\nu^c \quad (5)$$

where $\nu = \delta_c / \sigma$. Plot $b(z, M)$ versus M for $z = 0$ and $z = 1$ in the range $M = [10^{12}, 10^{16}]$.

Use log scale in the M axis and linear scale for $b(M)$.

b) For a fiducial cosmology, compute the halo profile for the fit of NFW:

$$\rho(r|M_{vir}, z) = \frac{\rho_s}{cr/r_{vir}(1 + cr/r_{vir})^2} \quad (6)$$

for a halo of mass $M_{vir} = 10^{14} M_\odot/h$ at redshift $z = 0$ and plot $\rho(r)$ versus r . You will need the formulae sequence below:

$$E^2(z) = \Omega_m(1+z)^3 + \Omega_{DE}(1+z)^{3(1+w)}, \quad (7)$$

$$\rho_{crit}(z) = \rho_{crit,0} E^2(z), \quad (8)$$

$$\omega_m(z) = \Omega_m(1+z)^3 / E^2(z), \quad (9)$$

$$x = \omega_m(z) - 1, \quad (10)$$

$$\Delta_c = 18\pi^2 + 82x - 39x^2, \quad (11)$$

$$\Delta_c = \frac{3M_{vir}}{\rho_{crit}(z)4\pi r_{vir}^3} \rightarrow \text{Find } r_{vir}, \quad (12)$$

$$\nu(M^*) = 1 \rightarrow \text{Find } M^*, \quad (13)$$

$$c(M_{vir}, z) = \frac{9}{1+z} \left[\frac{M_{vir}}{M^*} \right]^{-0.13} \rightarrow \text{Find } c, \quad (14)$$

$$M_{vir} = 4\pi\rho_s \frac{r_{vir}^3}{c^3} \left[\ln(1+c) - \frac{c}{1+c} \right] \rightarrow \text{Find } \rho_s, \quad (15)$$

2) Bonus Problem: Halo Model: Combine the Halo Model ingredients you found before to compute the Halo-Model power spectrum at $z = 0$ with the formulae sequence:

$$u(k|M) = \int_0^{r_{vir}} dr 4\pi r^2 \frac{\sin kr}{kr} \frac{\rho(r|M)}{M} \quad \text{see Cooray \& Sheth 2002} \quad (16)$$

$$P^{1h}(k) = \int d \ln M \frac{M^2}{\bar{\rho}_{m0}} \frac{dn}{d \ln M} |u(k|M)|^2 \quad (17)$$

$$P^{2h}(k) = \left[\int d \ln M \frac{M}{\bar{\rho}_{m0}} \frac{dn}{d \ln M} b(M) u(k|M) \right]^2 \quad P_L(k) \approx P_L(k) \text{ when } k \rightarrow 0 \quad (18)$$

$$P(k) = P^{1h}(k) + P^{2h}(k) \quad (19)$$

Plot $P_L(k)$, $P^{1h}(k)$, $P^{2h}(k)$ and $P(k)$, as well as the non-linear spectrum from `halofit`.